

Diseases of Turnip and Mustard Greens

Sherrie Smith
Plant Health Clinic
Diagnostician

Rick Cartwright
Professor - Plant
Pathologist

Introduction

Greens, the leaves of turnips, mustard, collards and related *Brassica* spp. (mustard family) plants, have long been a popular addition to the early fall garden. These plants are easy to grow but subject to several fungal and bacterial leaf spot diseases when environmental conditions are favorable for infection.

Common diseases of greens in Arkansas include:

Anthracnose (*Colletotrichum higginsianum*)

Cercospora leaf spot
(*Cercospora brassicicola*)

White spot (*Pseudocercospora capsellae*)

Downy mildew (*Peronospora parasitica*)

Bacterial leaf spot
(*Pseudomonas maculicola*)

Disease Development and Spread

These diseases are spread by wind or rain and are strongly favored by wet conditions. Anthracnose survives on volunteer plants, dead leaves and related weed hosts and is most severe during wet conditions with warm temperatures between 79°-86°F (warm, rainy periods during August and September). The pathogens that cause Cercospora leaf spot and white spot survive on volunteer plants and perennial weeds, and less commonly



Anthracnose symptoms include small, gray- to straw-colored spots on leaves.

Sherrie Smith

on infected seed. Cool temperatures of 55°-64°F and frequent rain, heavy dew or overhead irrigation increase both diseases. Night temperatures of 46°-61°F with daytime temperatures below 75°F favor downy mildew. The downy mildew fungus overwinters on *Brassica* spp. residue and is also more severe under frequent wet conditions. Bacterial leaf spot is favored by cool weather where daytime highs only reach 65°-75°F. The bacterial leaf spot pathogen can survive on crop residue for several years and on seed. Weather conditions in late September or during October and November in Arkansas favor all diseases except anthracnose (earlier disease).

Symptoms

Anthracnose attacks turnips, mustard, broccoli, collards and kale by producing small, gray- to straw-colored circular spots on the leaves and petioles. Spots on the stalks are sunken, elongated and gray to brown with a black border.

*Arkansas Is
Our Campus*

Visit our web site at:
<http://www.uaex.edu>

Symptoms of white spot include circular spots with gray, brown or nearly white centers on cotyledons, leaves and petioles. Edges of the lesions are slightly darkened. Leaves turn yellow and drop prematurely with this disease. Mustard, collards and turnip greens are all susceptible. Seedlings may be killed if the disease is severe at emergence.

Cercospora leaf spot lesions are pale green to gray or white and often have a brown border. Lesions are angular or circular in shape. This disease is sometimes called frog-eye leaf spot. Plants can be defoliated when infections are severe.

Downy mildew usually attacks lower leaves first. Small, yellow-brown spots appear which eventually expand and develop gray to black lace-like or streaky markings. In moist weather, white fungal mats are visible on the underside of leaf spots. Downy mildew can spread very fast under ideal (wet, cool) conditions.

Bacterial leaf spot symptoms begin on leaves as numerous, pinpoint-sized and irregular brown spots with yellow halos. This stage is difficult to see on the upper leaf surface, whereas water-soaked brown spots are clearly visible on the leaf underside. As the disease progresses, the many small spots coalesce into large, brown papery areas with yellow borders that tear, giving the plant a ragged appearance.

Disease Control

- Start with a soil test before planting and fertilize according to test results to ensure adequate, balanced fertility.
- Plant in warm, well-drained and tilled soil.
- Plant certified clean seed and healthy transplants. If certified seed is not available, use an approved seed treatment.
- Practice good weed control in and around the field. Remove all volunteer crucifer (*Brassica*)



David B. Langston, University of Georgia

Circular spots with gray, brown or nearly white centers on cotyledons, leaves and petioles are symptoms of white spot.



Ohio State University Extension

Visible white fungal mats on the underside of leaves are evidence of downy mildew.



Sherrie Smith

Cercospora leaf spot lesions are pale green to gray or white and angular in shape.



Sherrie Smith

Bacterial leaf spot progresses into large, brown papery areas with yellow borders giving the plant a ragged appearance.

spp.) weeds such as wild mustards, shepherd's purse, pepper-grasses and wild radish.

- Practice crop rotation and do not grow crucifers again for at least three years on the same spot.
- Monitor and use pesticides as necessary for insect and disease control (see table).
- Remove any diseased plants immediately, if possible, to prevent disease buildup and spread.
- Till all old crop debris immediately after harvest to allow time to rot.
- Avoid overhead sprinkler irrigation if possible as this makes disease worse. Work in the fields only when leaves are dry to prevent

spreading disease with your hands, tools or clothing.

References

Averre, C.W., Jenkins, S.F. 2006 online. Diseases of Greens and Their Control.

<http://www.growit.com/bin/Problems.exe?MyType=376>

Brassica Notes. 1989. D. Undersander. University of Wisconsin-Extension. FC 15.4.1.

Black, L.L. AVRDC. Vegetable Diseases: A Practical Guide. 2001.

Motes, J.E., Cartwright, B., Damicone, J.P. 2006 online. Greens, Production (Spinach, Turnip, Mustard, Collard, and Kale). <http://www.osuestra.com>

Shikora, E.J. 2006 online. Plant Disease Notes Leaf Spots of Mustard and Turnip. Alabama Cooperative Extension. www.aces.edu

Fungicides (0 = ineffective, 4 = very effective)

Commercial Production	Preharvest Interval (DAYS)	White Spot			
		Downy Mildew	Cercospora Leaf Spot	Anthraco	Bacterial Leaf Spot
azoxystrobin (Amistar, Quadris)	0	3	3	4	0
cyprodinil + fludioxonil (Switch)	7	0	2	1	0
fixed copper (Kocide)	0	2	1	1	3
maneb (Manex)	7	3	2	3	0
pyraclostrobin (Cabrio EG)	3	3	4	4	0
Home Gardens					
Copper hydroxide (Kocide)	0	2	1	1	3

Disclaimer: THE LABEL IS THE LAW. The above information should be used as a general guide on effectiveness, but actual use and other factors are strictly governed by the registered and latest label on the product. Always read and follow the label. This information is for educational purposes only and was as accurate as could be determined as of November 30, 2006. The University of Arkansas Division of Agriculture makes no purposeful distinctions between brands or products mentioned, except where scientific research has been published to indicate differences.

Printed by University of Arkansas Cooperative Extension Service Printing Services.

SHERRIE SMITH is plant health clinic diagnostician, University of Arkansas Division of Agriculture, Cooperative Extension Service, Lonoke. **DR. RICK CARTWRIGHT** is professor - plant pathologist, University of Arkansas Division of Agriculture, Cooperative Extension Service, Fayetteville.

FSA7549-PD-1-07N

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director, Cooperative Extension Service, University of Arkansas. The Arkansas Cooperative Extension Service offers its programs to all eligible persons regardless of race, color, national origin, religion, gender, age, disability, marital or veteran status, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.