

STORED GRAIN INSECT CONTROL

Type Treatment	Insect	Insecticide	Rate	Application/Comments
Preventive treatments for bins (empty bins, walls, ceilings, beams, outside storage, foundations and grain handling equipment)	Beetles and Moths	Storcide II	(Rate/1,000 sq ft) 1.8 oz/1 gal water	Clean bins before application. Storcide II may be applied using automated spray equipment. Do not enter bin until spray has dried. Use malathion for wooden or metal bins. Malathion will not control Indian meal moth. Malathion is NOT labeled for rice storage facilities. DO NOT APPLY DIRECTLY TO GRAIN.
		Malathion 5 EC premium grade (Fyfanon) Tempo Ultra SC	5.12 oz/gal water 0.54 oz/gal water	
Preventive treatment sprays for grain	Beetles and Moths	Actellic 5 E	9.2-12.3 oz/30 tons grain (1,071 bushels)	Shelled corn and grain sorghum
		Malathion 6% Dust	10 lb/1,000 bushels (wheat, corn, oats, rye or barley)	Apply as the grain is being loaded or moved into final storage for protection from beetles and moths. Malathion is not labeled for rice.
		Storcide II (Storcide II is not labeled for corn)	(Rate/1,000 bushels) Barley – 9.9 fl oz Oats – 6.6 fl oz Rice – 9.3 fl oz Grain Sorghum – 11.6 fl oz Wheat – 12.4 fl oz	Apply Storcide II in 1-5 gal of water into main grain stream as a coarse spray.
Surface treatment (inside bins and grain treatment)	Beetles and Moths	Malathion 6% Dust	5 lb/1,000 sq ft	Apply as top-dress on grain bin surface for protection against moths and beetles. Malathion is not labeled for rice.
	Moths only	<i>Bacillus thuringiensis</i> (Dipel, SOK-Bt, etc.)	Follow label directions.	Apply as top-dress for protection against moths only. Incorporate into top 4 inches.
Grain Fumigation	Beetles and Moths	Aluminum Phosphide (R) (Phostoxin, Fumitoxin, Phoskill, Phostek)	Silos and Concrete Bins 200-900 pellets/1,000 bu 40-180 tablets/1,000 bu 10 pellets/ton 6 tablets/ton (check label also) Butler-Type Bins 450-900 pellets/1,000 bu 90-180 tablets/1,000 bu Space Treatments pellets: 100-725/1,000 cubic ft tablets: 20-145/1,000 cubic ft	Grain surface should be level. Apply in gas-tight bins when grain temperature is 60°F or higher. Leave for at least 3 days, preferably 5 days. Do not fumigate when temperatures are below 40°F. Use higher rates when infestations are high and/or temperatures are low. Always use a gas mask and never fumigate alone. Fumigated grain should be aerated thoroughly before feeding. Use a self-contained breathing apparatus and a safety rope. Seal all buildings and storage bins airtight before fumigation. Fumigation is not recommended for storage bins not equipped with fans for aeration.
Grain Fumigation	Beetles and Moths	Aluminum Phosphide (R) (Phostoxin, Fumitoxin, Phoskill, Phostek)	Bagged Grain or Feed 150-450 pellets/1,000 cubic ft 30-90 tablets/1,000 cubic ft	Be careful under hot, humid conditions when using aluminum phosphide because phosphine gas may be released quickly under these conditions. Phosphine needs at least 72 hours to kill some immature insects. Special training and precautions are needed before using aluminum phos- phide. For more information on treatment using aluminum phosphide: http://www.degeschamerica.com/index.shtml .
HOME STORAGE Protection of small quantities of grain for home storage to be eaten	Beetles and Moths	Heat oven until center of grain is 120 degrees F.	30 minutes	Place in tight container. Temperature should be checked with meat thermometer. Do not use heat on seed to be planted.
		Cold or freezer treatment at 0 degrees F.	4 days	Place container in deep freeze.