

STRAWBERRY INSECT CONTROL

Pest	Material and Formulation	Amount to Use Per Acre	Remarks/Precautions	Days to Harvest
PREBLOOM				
Strawberry Clipper	Brigade WSB	6.4-32 oz	Begin checking for clipped buds when the first flower appears. If clipped buds are found, apply insecticide and repeat spray at 10-day intervals as long as bud clipping continues. If spittlebugs are a problem, spray when buds first become visible and make a second application just before the first blossom opens. Use higher rates of Danitol or Brigade for plant bug and spittlebug control. Lorsban applications are prebloom only.	0
	Danitol 2.4EC	10 2/3 oz		2
	Lorsban 4E	1 qt		21
Tarnished Plant Bug Spittlebug	Endosulfan EC	1 1/3 qt		14
	Sevin 80S	2.5 lb		7
	Thiodan	1 1/3 qt		14
	Danitol 2.4 EC	10.67 fl oz		2
	Provado 1.6	3.8 fl oz		7
Spider Mites	Abba 0.15 EC	16 fl oz	Make two applications of Abba at 7- to 10-day intervals. Do not retreat within 21 days of second application. Allow two sprays of Acramite per season. Do not make more than two applications of Kanemite per season. Danitol or Brigade at higher rates control spider mites. Road dust control can reduce mite buildup. Do not use the following with oil or within 2 weeks of applying oil: Dyrene, Bravo, Botran, Kelthane, Dikar, Karathane, Ambush or Pounce.	3
	Acramite 50 WS	0.75-1.0 lb		1
	Kanemite 15 SC	21-31 fl oz		1
	Brigade WSB	16-32 oz		0
	Vendex	2 lb		1
	Kelthane 35W	1-2 lb		3
	Danitol 2.4EC	10 2/3 oz		2
	Agri-Mek 0.15EC	16 fl oz		3
Cyclamen Mite	Savey	6 oz		3
	JMS Stylet-oil	1-2 gal		
	Thiodan EC	4 qt		4
	Endosulfan EC	2 2/3 qt		4
	Kelthane 35W	3-4 lb		3
EARLY BLOOM THROUGH BLOOM			SAVE THE BEES!! DO NOT APPLY INSECTICIDES DURING BLOOM.	
POST-BLOOM TO HARVEST				
Aphids, Whiteflies	Admire Pro	10.5-14 fl oz	Do not exceed 14 fl oz per acre per season.	14
Leafrollers	Javelin WG	0.25-1.5 lb	A <i>Bt</i> formulation approved for organic production.	0
POST-HARVEST				
White Grubs of Japanese Beetle	Admire Pro	7-10.5 fl oz	Surface application followed by 0.25 inch of rain or overhead irrigation. Do not exceed 14 fl oz per acre per season.	14
	Alias 2 F	16-24 fl oz	Surface application followed by 0.25 inch of rain or overhead irrigation. Do not exceed 24 fl oz per acre per season.	14
	Platinum 2SC	5-12 oz		50

RED IMPORTED FIRE ANTS

ALL FIRE ANT BAITS – Apply when ants are active and soil temperature is above 60 degrees F. Do not treat if rain is anticipated within 6 hours.

S - Methoprene (Extinguish)	3-5 T/mound 1-1.5 lb/A	Mound-to-mound treatment rate. Broadcast rate. This product is an insect growth regulator (IGR). Apply 3 to 4 weeks prior to harvest date of production area being treated.
Pyriproxyfen (Esteem)	2-4 T/mound 1.5-2.0 lb/A	Mound-to-mound treatment rate. Broadcast rate.