

CANOLA DISEASES

Rick Cartwright and Scott Monfort

Disease	Fungicide	Active Ingredient	Rate/Acre	Days to Harvest	Comments
Blackleg (<i>Leptosphaeria</i>)	Quadris	azoxystrobin	6 - 15.5 fl oz	30	Group 11 fungicide. See label.
Seedborne Blackleg Seedling Rhizoctonia Damping Off Seedling Blight (<i>Alternaria</i>)	Dynasty	azoxystrobin	0.1 - 3.75 fl oz/cwt (seed treatment)	See label.	Group 11 fungicide. See label.
Sclerotinia Stem Rot	Endura	boscalid	5 - 6 fl oz	21	Maximum of 2 applications per season. Group 7 fungicide. Apply at 20% - 50% bloom stage.
	Proline 480SC	prothioconazole	4.3 - 5.7 fl oz	36	Maximum of 2 applications per year. Apply at 20% - 50% bloom stage. Group 3 fungicide.
	Quadris	azoxystrobin	6 - 15.5 fl oz	30	Group 11 fungicide. See label.

This information was current as of October 1, 2009, and applies only to Arkansas and may not be appropriate for other states or locations. The listing of any product in this publication does not imply endorsement of that product or discrimination against any other product by the University of Arkansas Division of Agriculture. *Every effort was made to ensure accuracy, but the user of any crop protection product must read and follow the most current label on the product – The Label is the Law.* For further assistance, contact the local Cooperative Extension Service office.