

## HOME LAWN DISEASES

Stephen Vann

Disease	Product	Active Ingredient	Rate	Comments
Brown Patch, Dollar Spot, Rusts, Leaf Spots	Scotts Lawn Fungus Control	thiophanate-methyl	1.34 lb/1,000 sq ft	Apply at first sign of disease. Irrigate following application. See label. Use lawn spreader for application.
	Green Light Systemic Fungicide	thiophanate-methyl	2 oz/1,000 sq ft	See label.
	Fertilome Liquid Systemic Fungicide	propiconazole	6 - 10 tbs	Apply at first sign of disease using 1 gal spray per 250 sq ft (10' x 25').
	Maxide Disease Killer	azoxystrobin	2 - 4 lb/1,000 sq ft	14 - 21 day application interval.
	Fertilome Halt Systemic Rose, Flower, Lawn and Ornamental Fungicide	thiophanate-methyl	2 oz/1,000 sq ft	3 - 5 gals spray mix/1,000 sq ft. 7 - 14 day interval.
	Ortho Lawn Disease Control	propiconazole	See label.	Packaged as hose-end applicator.
	Monterey Fungi-Fighter	propiconazole	3 - 5 oz/250 sq ft	See label.
	Spectracide Immunox (liquid)	myclobutanil	7 tbs	Apply at first sign of disease using 1 gal spray per 250 sq ft (10' x 25'). Use higher rate for brown patch or severe disease. Apply at 10 - 14 day intervals as needed.
	Spectracide Immunox Lawn Dis. Control (granules)	myclobutanil	See label.	Apply at first sign of disease.
	Green Light Fung-Away Systemic Granules	myclobutanil	3 - 6 lb/1,000 sq ft	Apply at first sign of disease. See label.
	Green Light Fung-Away Systemic Lawn Spray	triadimefon	3 - 6 oz/1 gal/100 sq ft	See label.
	Natural Guard Copper Soap Liquid	copper octanoate	1.5 - 6 oz/2.5 gal/ 1,000 sq ft	See label.
	Fertilome F-Stop Granules	myclobutanil	4 - 8 lb/1,000 sq ft	Apply at first sign of disease. See label.
	Hi-Yield PCNB Granular Fungicide	PCNB	See label.	Apply at first evidence of disease activity. Apply with lawn spreader.
	Hi-Yield Lawn Fungicide Granules containing Bayleton	triadimefon	3 - 6 lb/1,000 sq ft	Apply to dry grass at first sign of disease and again at 15 - 30 day intervals if needed. Uniform application is critical. Sprinkle-irrigate to activate after application – at least 1/4" rain or irrigation is necessary.
Hi-Yield Turf and Ornamental Fungicide containing 10% PCNB	PCNB	5 - 7.5 lb/1,000 sq ft	See label for specific rates. Irrigate after application.	
Bayer Advanced Fungus Control for Lawns	triadimefon	See label.	Apply to dry grass at first sign of disease. See label.	
Bonide Fung-onil Lawn Disease Control	triadimefon	3 - 6 lb/1,000 sq ft	See label.	
Hi-Yield Captan 50WP	captan	5 tsp/gal/100 sq ft	7 - 10 day interval – begin with new growth in spring. Do not exceed 2 applications per year.	
H & I Agritech GreenCure	potassium bicarbonate	1 - 2 tbs/gal/450 sq ft	Begin at first sign of disease. 7 - 14 day intervals. For organic production.	
Pythium Diseases	Monterey Aliette	aluminum tris	4 oz	Spray onto foliage in 1 - 5 gals water per 1,000 sq ft.

**CAUTION: Chlorothalonil is prohibited on home lawns (turf).**