

Feed, Forage and Grazing Restrictions for Cotton Herbicides

Herbicide	Restrictions
Aim	Do not graze treated fields or harvest for forage or hay.
Assure II	Do not graze treated fields or harvest for forage or hay.
Buctril	Do not graze any portion of crop. Do not cut for feed or fodder.
Caparol	Do not feed treated forage to livestock or graze treated areas or illegal residues may result.
Cobra	Do not graze animals on green forage or stubble. Do not utilize hay or straw for feed or bedding.
Direx	Do not allow livestock to graze treated corn.
DSMA	Do not feed treated foliage to livestock or graze treated areas.
Dual Magnum	No information on label.
Envoke	Do not graze or harvest for forage or hay.
Fusilade DX	Do not graze or harvest for forage or hay.
Glyphosate	Do not graze or feed.
Ignite	No information on label.
Linex	Do not graze treated fields or feed forage from treated areas to livestock. Do not feed gin trash to livestock
Meturon	Do not feed foliage from treated cotton plants or gin trash to livestock.
MSMA	Do not feed treated foliage to livestock or graze treated areas.
Poast Plus	Do not graze treated cotton fields, and do not feed treated cotton forage to livestock.
Prowl	Do not feed forage or graze livestock in treated cotton fields.
Reflex	Do not graze or harvest for forage or hay.
Select	Do not graze treated fields or feed treated forage or hay to livestock.
Staple LX	Do not feed treated gin by-products (trash) to livestock.
Treflan	No information on label.
Valor	Do not graze or harvest for forage or hay.

Restrictions are listed as worded on the labels. Feeding and application restrictions for herbicides are generally based on residue tolerances allowed for animal feeding. The restrictions are generally not due to acute toxicity (poisoning) problems. Livestock that are accidentally fed treated crops earlier than allowed may not be harmed but may have illegal pesticide residues in their meat or milk. If you have fed livestock treated crops within the restricted period, refer to the label, your dealer, or herbicide company representative for more information.

*Many labels refer to soybean plants as 'vines'.

Cotton Postemergence Herbicide Preharvest Application Intervals (PHI)

Herbicide	PHI
Aim	7 days
Assure II	80 days
Buctril	60 days
Caparol/Cotton Pro	No restrictions.
Cobra	70 days
Direx	No restrictions.
DSMA	1st bloom
Envoke	60 days
Fusilade DX	90 days
Glyphosate	7 days
Ignite	70 days
Linex	No restrictions.
Meturon	60 days
MSMA	1st bloom
Poast Plus	40 days
Select	60 days
Staple	60 days
Valor LX	60 days

These intervals are the number of days that must be allowed between herbicide application and harvest. Applications made after these interval restrictions could cause illegal herbicide residues to be present in the harvested seed or fiber.