

Crop Replant and Rotation Guide for Soybean Herbicides*

Herbicide	Replant/ Crop Rotation	Time Interval	Precautions
Assure II	S,C All	I 120 days	
Authority MTZ	S B,W,SG A,CT,GS,P,SF	I 4 months 12 months	
Basagran	All	I	
Butyrac/Butoxone (2,4-DB)			No restrictions.
Cadet	All	I	
Canopy	S W,B,RG C AL,CT,GS,R CL P All	I 4 months 9 months 10 months 12 months 8 months 18 months†	Use only on soils with pH of 7.0 or less. †Successful field bioassay must be completed prior to planting. Wait 15 months before trying a bioassay.
Canopy (pH 7.1 to 7.5)	S W,B,RG AL,CL,C, CT,GS,R,P All others	I 4 months 18 months 18 months†	Use only in soils with pH 7.5 or less. †Successful field bioassay must be completed prior to planting. Wait 15 months before trying a bioassay.
Canopy EX	W C,GS,R,AL,CL CT SF,CA	3 months 9 months 10 months 18 months	See label for pH, rate and geographical restrictions. High pH soils have longer rotational intervals.
Classic	S† SG,RG P C,CT CL,GS,R All	I 3 months 6 months 8 months 18 months††	If applied after Aug. 1, extend recrop interval for C, CT, R and GS 2 months. †Recrop intervals may also be extended if Classic is applied after application of Canopy. See labels for information. ††Successful field bioassay must be completed prior to planting. Wait 9 months before trying a bioassay.
Cobra			No restrictions.
Dual Magnum or Sequence	S,C,CT,GS† SG,W Rice All	I 4.5 months Next spring 18 months	†Use Concep-treated grain sorghum seed.

Herbicide	Replant/ Crop Rotation	Time Interval	Precautions
Enlight	W CT,C,R,SF AL,CL	4 months 9 months 12 months	
Envive	W,CT C,GS,R,AL,CL SF,CA	4 months 10 months 18 months	See label for pH and geographical restrictions. High pH soils have longer rotational intervals.
FirstRate	W C,CT,P,R,GS	3 months 9 months	All others – 30 months plus bioassay.
Flexstar GT	CT,S W C,P,R,GS Other	0 months 4 months 10 months 18 months	See Reflex.
Fusilade DX or Fusion	S, CT All	I 2 months	
Glyphosate	All	I	
Ignite	C,S,CT,R,CA W,B,O GS	I 70 days 180 days	No restrictions.
Outlook	C,S SG All	I 4 months FY	Do not retreat field with second application.
Paraquat			No restrictions.
Poast/Poast Plus			See label.
Prefix	S CT AL,W,O,R,RY,B GS,R,C	I 1 month 4.5 months 10 months	Use Concep-treated grain sorghum seed. Do not use Reflex or Flexstar in-season.
Prowl	S,CT W,B All	I 4 months† FY	Do not rework soil deeper than treated zone. † Cannot replant using no-tillage practices.
Pursuit	S B,W C Rice All	I† 4 months 9.5 months 40 months 18 months	†Do not rework soil deeper than 2 inches. Do not apply Classic, Canopy, Lorox Plus, Scepter, Squadron or Tri-Scept the same year as Pursuit or injury to following crops may occur.

NOTE: See page 38 for key to abbreviations.

(continued on page 38)

Crop Replant and Rotation Guide for Soybean Herbicides* [cont.]

Herbicide	Replant/ Crop Rotation	Time Interval	Precautions
Python	AL,P,B,W,RY, O R C GS Sunflower CT All	4 months 6 months I 12 months 18 months 18 months 26 months	Requires successful bioassay.
Reflex or Flexstar	S SG, W C,CT,R,P,GS All	I 4 months 10 months 18 months	
Resource	SG S,C All	120 days I 30 days	
Scepter	S R W B,C* GS†,O,RY P All	I FY 4 months 11 months 18 months	Do not apply Classic, Canopy, Lorox Plus, Pursuit or Pursuit Plus the same year as Scepter or injury to following crops may occur. * Do not plant corn or grain sorghum for 18 months if Scepter was used as a sequential spray pre (or ppi) and post. Do not plant corn the next year if rainfall was less than 15 inches from the time of application through October (or within 6 months of application in the Boot-heel.) If replanting soybeans following crop loss, do not rework soil deeper than 2 inches.
Select Max	C		See label.
Sencor	S† AL,C,FG,W,B CT,R All (except root crops)†† Root crops	I 4 months 8 months 12 months 18 months	† Waiting period for replanting soybeans depends on the rate of metribuzin used. See specific label for more information. Add 2 months to time intervals if pH of soil is above 7.5. †† Cover crops may be planted anytime, but stand reductions may occur.

Herbicide	Replant/ Crop Rotation	Time Interval	Precautions
Storm	All	Fall	
Treflan/others	S,CT W,B All	I Fall FY	
Treflan (2X rate)	S,CT R All	I FY 2 years	
Typhoon	W C,CT,R GS & all others	4 months 10 months 18 months	
Ultra Blazer	S All	I Fall	
Valor	S,P W,C,R,GS CT B A,O,CL,CA	I 30 days 21 days 4 months 8 months	Pre-emerge treatments may injure soybeans.

Key		
Crop		
All = All crops not specified	FG = Forage Grasses	RG = Ryegrass
AL = Alfalfa	FL = Forage legumes	RY = Rye
B = Barley	GS = Grain Sorghum	S = Soybeans
C = Corn	O = Oats	SF = Sunflowers
CA = Canola	P = Peanuts	SG = Small Grains
CL = Clovers	R = Rice	W = Wheat
CT = Cotton		
Timing		
I = Immediately		
FY = Following year (usually spring or following fall - 11 to 16 months)		

*This table applies to the major field and forage crops. Refer to the herbicide labels for the latest recrop and rotation information for horticultural crops. These are written as best we could interpret the labels. We regret any omissions or errors. Always refer to product labels before using a pesticide or replanting into treated fields.