

**Forage, Feed and Grazing Restrictions  
for Soybean Herbicides**

Herbicide	Restrictions
Assure II	Do not graze treated fields or harvest for forage or hay. Do not apply within 80 days of harvest.
Basagran	No restrictions on label.
Broadstrike + Dual or Treflan, Python	Do not graze or feed treated soybean forage, hay or straw to livestock.
Butyrac/Butoxone (2,4-DB)	Do not graze or feed soybean hay within 60 days after application of a 2,4-DB tank-mix application. Do not harvest soybeans within 60 days after spray application.
Canopy	Do not graze treated fields or harvest for forage or hay.
Classic	Do not graze treated fields or harvest for forage or hay. Do not apply later than 60 days before soybean maturity.
Cobra	Do not graze animals on green forage or stubble. Do not utilize hay or straw for animal feed or bedding. Apply not later than 90 days before harvest.
Dual Magnum	No restrictions on label.
FirstRate	Do not harvest for forage or hay for 14 days.
Fusilade DX, Fusion	Do not graze or harvest for forage or hay.
Glyphosate	Do not harvest or feed treated crops for 8 weeks after application. Allow 14 days following spot treatment or selective equipment before grazing domestic livestock.
Gramoxone Max/Inteon	Do not graze treated areas or feed treated forage to livestock.
Ignite	Do not feed green soybean plants to livestock.
Outlook	Do not graze or feed forage, hay or straw to livestock.

Herbicide	Restrictions
Poast/ Poast Plus	Do not graze treated fields and do not feed treated soybean forage (green succulent) or ensilage to livestock. Treated soybean hay may be fed. Do not apply to soybeans within 90 days of harvest.
Prowl	Livestock <i>can</i> graze or be fed soybean forage from treated fields.
Pursuit	Do not graze or feed treated soybean forage, hay or straw to livestock. There should be an interval of at least 85 days between an application of Pursuit and soybean harvest.
Python	Do not graze or harvest for forage or hay.
Reflex/Flexstar	Do not graze treated areas or harvest for forage or hay. Do not graze rotated small grain crops or harvest for livestock forage or straw. Do not apply after first bloom.
Resource	Do not graze or harvest for forage or hay.
Scepter	Do not graze or feed treated soybean forage, hay or straw to livestock.
Select	Do not graze treated fields or feed treated forage or hay to livestock.
Sencor	Treated vines* may be grazed or fed to livestock 40 days after application.
Storm	Do not use treated plants for feed or forage. Do not apply within 50 days of harvest.
Synchrony	Do not graze treated fields or harvest for forage or hay. Do not apply later than 60 days before soybean maturity.
Treflan	No restrictions on label.
Typhoon	Do not graze or harvest for forage or hay.
Ultra Blazer	Do not use treated plants for feed or forage.
Valor	Do not graze or harvest for forage or hay.

Restrictions are listed as worded on the labels. Feeding and application restrictions for herbicides are generally based on residue tolerances allowed for animal feeding. The restrictions are generally not due to acute toxicity (poisoning) problems. Livestock that are accidentally fed treated crops earlier than allowed will not be harmed, but may have illegal pesticide residues in their meat or milk. If you have fed livestock treated crops within the restricted period, refer to the label, your dealer, or herbicide company representative for more information.

\* Many labels refer to soybean plants as 'vines'.