

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
Woody Plant Control in Non-Cropland (Right-of-ways, fencerows, industrial sites, etc.) Foliar Spray				
imazapyr @ 0.5 to 1.5 lb/A	Hickory, honeylocust, honeysuckle, kudzu, maple, mulberry, multiflora, rose, oaks, poison ivy, sassafras, sumac, sweetgum, willow, other broadleaf plants.	Arsenal 2S 2 to 6 pt of 2 lb/gal.	Apply to actively growing vegetation not under stress.	Apply during warm weather after full leaf out and before leaf drop. Apply to foliage in 10 to 60 gal/A if using ground equipment. Use 0.5 to 1% for low volume hand application. Spray to wet but do not allow runoff. Keep away from foliage or roots of desirable plants.
dicamba + 2,4-D amine 3 + 5.7 lb/A or dicamba + 2,4-D amine 0.125 + 0.35 to 0.5 + 1 lb/A	Persimmon, poison ivy, other broadleaf plants.	Weedmaster or (1 pt to 2 qt/A).	Apply uniformly as a foliar spray after leaves are fully developed until three weeks before a frost.	Wet foliage to point of runoff with a backpack sprayer. Apply in 200 to 300 gal per acre with a hydraulic sprayer. See label for application directions for specific brush types.
metsulfuron @ 0.038 to 0.15 lb/A	Kudzu, honeysuckle, black locust, sericea lespedeza, privet, bodark.	Escort 60 DF 1 to 4 oz.	During periods of active growth.	Use the 4 oz rate for kudzu and add 0.25% nonionic surfactant. Wet the kudzu canopy thoroughly using at least 60 gpa. Make a follow up treatment one year later to control escapes and misses. Failure to do so will result in reinfestation.
triclopyr @ 2.25 to 4 lb/A	Blackberry, oaks, pines, sumac, sweetgum and other broadleaf plants.	Garlon 3A or 4 0.75 to 1 gal.	Apply uniformly as a foliar spray after leaves are fully developed until three weeks before a frost.	Wet foliage to point of runoff with a backpack sprayer. Apply in 100 to 400 gal per acre with a hydraulic sprayer. See label for application directions for specific brush species.
bromacil @ 4.4 to 24 lb/A	Controls many brush and tree species.	Hyvar X-L 2.25 to 12 gal/A.	Apply to actively growing brush.	Do not apply to brush standing in water, do not use in irrigation ditches nor on right of ways or other sites where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated areas.
fosamine @ 6 to 12 lb/A	Kudzu, pines and other woody plants.	Krenite S + Surfactant (nonionic) 1.5 to 3 gal/A + 1 qt/100 gal of water.	Apply as a foliar spray from July until the first frost in the fall.	Complete coverage is required for control. See label for application directions for specific brush species.
glyphosate @ 1.5 to 3.75 lb/A	Most annual weeds and many perennials such as johnsongrass, bermudagrass, curly dock, milkweed, horsenettle, honeysuckle, lespedeza, brambles, multiflora rose and trumpet creeper.	Roundup Pro 4L + Surfactant (nonionic) 2 to 5 qt/A + 2 qt/100 gal of water.	When plants are actively growing. Perennial plants are best controlled when sprayed at growth stages near maturity. Brush control is best when applications are made in fall to non-stressed trees.	Follow instructions and precautions listed on the label. Adding 0.5% nonionic surfactant has improved control of some perennials. Surfactant must be added to Accord. Tank mix with Arsenal improves perennial grass control.
picloram + 2,4-D amine @ 0.54 + 2 lb to 2.2 + 8 lb/A	Dogwood, honeylocust, honeysuckle, persimmon, pines and other woody plants.	Tordon 101 Mixture 1 to 4 gal/A.	Apply to actively growing plants.	Use 15 to 25 gpa spray mix. See label for use rates for specific woody plant species. Restricted use pesticide.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
Woody Plant Control in Non-Cropland (Right-of-ways, fencerows, industrial sites, etc.)				
Soil Treatments				
bromacil @ 4.5 to 24 lb/A	Many woody plant species.	Hyvar X-L 2.25 to 12 gal/A.	Apply before or during the period of active growth when rainfall can be expected for activation.	Use higher rates (greater than 5 gpa) on high organic soils. Use as a soil treatment or basal treatment. For use on drainage ditches, use only as basal treatment. Apply broadcast treatment. Apply broadcast treatments using at least 200 gal per acre of water. Basal treatment may be applied undiluted using a hand-gun applicator, or mixed with water in a ratio of 1 gal Hyvar in 5 gal of water. Do not apply near desirable vegetation. See label for use roles for specific woody plant species.
tebuthiuron @ 2 to 6 lb/A	Dogwood, elm, honeylocust, honeysuckle, multiflora rose, oaks, sumac, and other woody plants.	Spike 20P or 80W 10 to 30 or 2.5 to 7.5 lb/A.	Before or during the period of active growth. Best applied in mid-March.	Apply in 15 to 150 gal of water per acre before or during the period of active growth of target plants. See label for use rates for specific plant species to be controlled. Do not broadcast where maintenance of a grass cover is desired. Has some postemergence activity on some herbaceous weeds. May be used as an individual plant treatment on forage or pasture area when used at less than 5 lb/A. Do not cut for hay for 1 year after application.
hexazinone @ 2 to 12 lb	Honeylocust, multiflora rose, oaks, and other woody plants.	Velpar L 1 to 6 gal.	Apply in late winter or early spring before rainfall that is needed for activation.	Direct spray to the soil beneath woody plants to be controlled.
Cut Surface (frill, injection, hypo-hatchet, stump)				
2,4-D amine (4 lb/gal formulation)	Elm, poplar, sassafras, willow and many woody species.	2,4-D amine (4 lb/gal formulation) Undiluted. 1 to 2 ml of concentrate per injection.	May to October.	Make injections as near the root collar as possible. See label for instructions for specific woody plant species.
imazapyr	Many woody plant species.	Arsenal 2S 2 qt per 1 qt of water (concentrated), or 8 to 12 fl oz per gal of water (dilute).	During active growth.	May be used as a cut stump, injection, frill, or girdle treatments. See label for instructions for specific uses and rates.
imazapyr	Many woody plant species.	Chopper RTU Undiluted.	During active growth.	Spray or brush the undiluted solution onto the cambium area of freshly cut stump surface and the bark of the cut stump. Insure that the cambium area (wood just inside the bark) is thoroughly wetted, but not to the point of puddling.
triclopyr	Maple, oaks, poison ivy, sumac, sweetgum, and other woody plant species.	Garlon 3A, Tahoe 3A Undiluted.	Apply in spring or summer.	Apply in 1 ml of concentrate to cuts spaced 3 inches apart around the tree trunk. May also be applied with frill or girdle method.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
glyphosate	Oaks, pines, poison ivy, sweetgum, sycamore, and other woody plant species.	Glyphosate (4 lb/gal formulations) See table on page 23 for specific glyphosate conversions and surfactants. Undiluted.	Apply during active growth.	Apply 1 ml in cuts spaced 2 to 3 inches apart around the tree trunk.
picloram + 2,4-D amine	Pines and other woody plant species.	Undiluted. Pathway	Any time except during heavy sap flow.	Apply .5 ml of undiluted solution to cuts spaced 3 inches apart around the trunk. Or use 1 ml of 50% solution of Tordon 101 in a continuous cut girdling the trunk. Use undiluted Tordon RTU for frill method. Restricted use pesticide.
Basal Spray imazapyr (see label)	Many brush species.	Stalker 2L (See label.)	Spring or Fall.	For thinline and low volume basal bark treatments. See label for specifics. Imazapyr is soil active and may be taken up by the roots of desirable vegetation.
2,4-D ester + triclopyr	Many woody plant species.	Crossbow 4 gal in enough diesel oil, No. 1 or No. 2 fuel oil, or kerosene to make 100 gal of spray mixture.	Apply anytime except when snow or water prevent spraying the groundline.	Spray basal parts of brush or trees to a height of 15 to 20 inches from the ground. Thoroughly wet all the basal bark area including crown buds and ground sprouts. Best results have been obtained with winter to early spring applications.
triclopyr	Blackberry, dogwood, hickory, maple, oaks, pines, sumac, sweetgum, and other woody plant species.	Garlon 4, Tahoe 4E 4 gal in enough Arborchem Basal Oil, diesel fuel, No. 1 or No. 2 fuel oil, or kerosene to make 100 gal of spray mixture.	Apply anytime except when snow or water prevent spraying the groundline.	For control of weedy plants with stems less than 6 inches in diameter. Spray the basal parts of trunks to a height of 12 to 15 inches from the ground. May be mixed in oil-water mixtures as well. Refer to label for rates and directions.
triclopyr @ 1.0 lb/gal	Many woody plants.	Pathfinder II Ready to use.	Apply anytime except when snow or water prevent spraying the groundline.	Use on plant with basal stem diameter less than 6 inches. Thoroughly wet the lower 12 to 15 inches of stems including the root collar, but not to runoff. See label for streamline basal bark instructions.
clopyralid @ 0.5 lb	Kudzu, honeylocust, black locust, sericea lespedeza.	Transline 3 lb/gal 1.33 pts.	Late June to early October.	Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions has the potential to contaminate groundwater which may be used for irrigation and drinking purposes. Users are advised not to apply picloram where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local county Extension office can provide further information of the type of soil in your area and the location of groundwater.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
General Herbaceous Weed Control in Non-Cropland (Right-of-ways, fencerows, industrial sites, etc.)				
imazapyr @ 0.5 to 3 lb/A	Ash, dogwood, greenbrier, hawthorne, and many other grass, broadleaf and brush weeds.	Arsenal 2S 2 to 6 pt/A.	May be applied preemergence or as postemergence spray.	Post sprays are usually more effective. Apply to wet foliage. Apply to actively growing vegetation.
2,4-D amine 1 to 4 lb/A	Many annual and perennial broadleaf weeds.	2,4-D amine 1 to 4 qt/A.	Postemergence.	Apply as a foliar spray in 15 to 30 gpa of water to young, vigorously growing weeds. Avoid drift to susceptible crops or other desirable vegetation.
dicamba @ 0.5 to 8 lb/A	Many annual and perennial broadleaf weeds.	Vanquish 4S 1 pt to 2 gal.	Postemergence.	Apply to actively growing weeds and brush. May be tank mixed with 2,4-D, Karmex, Dalapon, Princep, Tordon, Amitrole, Hyvar, Velpar, Spike, Garlon, and other herbicides to broaden spectrum of weed and brush control. See label for more information.
dicamba + diflufenzopyr @ 0.14 to 0.28 + 0.053 to 0.105 lb/A	Many broadleaf weeds.	Overdrive 76 DF 4 to 8 oz/A.	Postemergence.	Add 1 qt nonionic surfactant per 100 gallons of water. May be tank mixed with most pasture herbicides.
prodiamine @ 0.65 to 1.5 lb/A	Annual grasses.	Endurance 65 DF 1 to 2.3 lb/A.	Preemergence.	Tank mix with Roundup Pro or Arsenal for control of emerged weeds.
DSMA or MSMA @ 2.7 to 5.4 lb/A	Johnsongrass control in other perennial grasses.	DSMA or MSMA Many formulations.	Postemergence.	Apply when johnsongrass is 6 inches tall until early head stage. Repeat applications may be needed.
sulfosulfuron @ 0.062 to 0.13 lb/A	Johnsongrass, nutsedge, buttercup, others.	Outrider 75 DF 1.33 to 2.66 oz.	Postemergence to actively growing weeds. Treat johnsongrass at 12 to 18 inches.	Excellent for johnsongrass control in bermudagrass. Does not injure actively growing bermudagrass. Add 0.5% nonionic surfactant or methylated seed oil. May be tank mixed with Roundup Pro, Escort, Oust or other herbicides to broaden the spectrum of control.
triclopyr @ 1.0 to 4.5 lb/A	Many annual and perennial broadleaf weeds.	Tahoe 3A, Garlon 3A or 4 + Nonionic Surfactant 0.33 to 1.5 gal/A of Garlon 3A, or 1 to 4 qt/A of Garlon 4 + 0.25 to 1 pt per 20 to 100 gal of water.	Postemergence.	Apply any time during growing season. May be tank mixed with 2,4-D, or Tordon 22K to broaden spectrum of weed and brush control. See labels for more information.
fosamine @ 6 to 12 lb/A	Blackberry, multiflora rose, sumac, and other brush and woody plant species.	Krenite S + Nonionic Surfactant 1.5 to 3 gal/A.	Postemergence.	Apply as a foliar spray from July through first frost. Complete coverage is essential for good control.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
sulfometuron-methyl @ 0.5 to 0.56 lb/A	Johnsongrass, fescue, most annual grass and broadleaf weeds.	Oust 75 DF 1 to 12 oz/A.	Preemergence or postemergence.	Apply preemergence or early postemergence in late spring to early summer. Use nonionic surfactant for postemergence applications. Do not apply where runoff water may flow onto agricultural land or where other desirable vegetation is growing. May be tank-mixed with Karmex, Velpar and other herbicides for broader spectrum weed control. See label for more information.
sulfometuron + chlorsulfuron @ 0.14 to 0.048 to 0.35 + 0.12 lb/a	Many broadleaf weeds and grasses.	Landmark II MP 4 to 10 oz/A.	Preemergence and postemergence.	Premix of Oust + Telar.
glyphosate @ 0.75 to 3.75 lb/A	Johnsongrass, bermudagrass, fescue, dandelion, multiflora rose, thistles, most annual weeds, and many perennial plants.	Glyphosate (4 lb/gal formulations) See table on page 23 for specific glyphosate conversions and surfactants. 2 to 5 qt/A.	Postemergence.	Apply as foliar spray to actively growing plants. See label for use rates for specific plant species.
imazapic @ 0.032 to 0.188 lb/A	Johnsongrass, ragweed, tall fescue, prickly sida, trumpet-creeper.	Plateau 2S 2 oz to 12 oz/A.	Postemergence.	Provides weed control and growth suppression.
imazapic + glyphosate @ 0.023 + 0.063 to 0.188 + 0.5 lb/A	Johnsongrass, crabgrass, sandbur, sedges, tall fescue control or seedhead suppression.	Journey 4 to 32 oz/A.	Postemergence.	Do not use unless bermudagrass injury can be tolerated.
pendimethalin @ 2 to 4 lb/A	Annual grasses.	Pendulum 60 DF 3.3 to 6.6 lb/A.	Preemergence.	Tank mix with Roundup Pro or Arsenal for control of emerged weeds.
Soil Sterilants (Right-of-ways, fencerows, industrial sites, etc.)				
bromacil + diuron @ 1.6 + 1.6 to 18.4 + 18.4 lb/A	Many annual and perennial broadleaf and grass weeds.	Krovor II DF 2 to 23 lb/A.	Preemergence or early postemergence.	Apply just before weed emergence or in early stages of weed growth. See label for use rates for specific weeds.
prometon @ 8 to 20 lb/A	Many annual and perennial broadleaf and grass weeds.	Pramitol 25E 4 to 10 gal/A.	Preemergence or postemergence.	Apply prior to emergence until 3 months after weed emergence. Will give residual control for over 1 year. See label for use rates for specific weeds and uses.
prometon + simazine + sodium chlorate + sodium metaborate	Many annual and perennial broadleaf and grass weeds.	Pramitol 5 PS 0.35 to 0.92 lb/100 sq ft.	Preemergence or postemergence.	Apply prior to emergence until 3 months after weed emergence. Will give residual control for over 1 year. See label for use rates for specific weeds and uses.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
Soil Sterilants (Right-of-ways, fencerows, industrial sites, etc.) [cont.]				
tebuthiuron @ 1 to 4 lb/A	Many annual and perennial broadleaf and grass weeds.	Spike 20P 10 to 20 lb/A of 20P.	March.	May be applied any time except when ground is frozen or the soil is saturated with moisture. Do not apply near desirable vegetation where roots may come in contact with the herbicide. Avoid contamination of irrigation water. See label for use rates for specific weeds.
dichlobenil @ 4 to 8 lb/A	Many annual and perennial weeds and woody plant species.	Casoron 4G 100 to 200 lb/A.	Preemergence.	See label for application instructions, use rates and specific weeds.
bromacil @ 3 to 24 lb/A	Many annual and perennial broadleaf and grass weeds.	Hyvar X-L 1.5 to 12 gal/A.	Preemergence or postemergence.	Apply as spray in 100 to 200 gal of water per acre. See label for use rates for specific weeds.
diuron @ 4 to 12 lb/A	Many annual and perennial broadleaf and grass weeds.	Karmex 80DF 5 to 15 lb/A.	Preemergence or early postemergence.	Apply to soil shortly before weed growth begins. See label for use rates for specific weeds.
bromacil + diuron @ 1.6 + 1.6 to 12 + 12 lb/A	Many annual and perennial broadleaf and grass weeds.	Krovar IDF 4 to 30 lb/A.	Preemergence or early postemergence.	Apply just before weed emergence or in early stages of weed growth. See label for use rates for specific weeds.
hexazinone @ 6 to 12 lb/A	Many annual and perennial broadleaf and grass weeds.	Velpar L 3 to 6 gal/A.	Preemergence or early postemergence.	Apply to soil from late winter to early summer, or in fall. Needs rainfall for activation. See label for use rates for specific weeds.
imazapyr + diuron @ 1.0 + 8.0 to 1.5 + 12.0 lb/A	Most herbaceous weeds.	Sahara DG 13-19 lb/A.	Pre or postemergence.	Tank mix with Roundup Pro for faster burndown of emerged vegetation. Do not apply over or near the roots of desirable trees.