

RICE INSECTICIDE PERFORMANCE RATING, 2012

Insecticide	Restricted Use (R)	Chinch Bug	Fall Armyworm	True Armyworm	Short Horned Grasshopper	Rice Stink Bug		Rice Water Weevil (adult)	Rice Water Weevil (egg)	Midge Larvae	Rice Stalk Borer	Aphids (Greenbug, Oat Bird, Cherry Aphid)	Grape Colaspis
						Adults	Eggs and Immatures						
Cruiser		6		2				6	7				8
Demacor X-100	X	1		7				1	8		8		4
Dimilin 2L	X							0	7			0	
Karate Z	X	7	8	8	7	6	8	9		6	5	8	
Malathion		1	2	5	6	5	5	0				4	
Methyl Parathion	X	6	6	8	6	9	6	0				7	
Mustang Max	X	7	8	8	7	6	8	9			5	8	
Prolex/Proaxis/Declare	X	7	8	8	7	6	8	9			5	8	
Sevin		6	6	5	6	7	5					1	

Rating Scale: 0 = no control, 10 = excellent

The performance ratings in the chart are for comparison purposes only and are not necessarily a measure of percent control.

Shaded boxes indicate products recommended for specific pests in this guide.

QUICK VIEW THRESHOLD GUIDE

Insect	Threshold				Scouting Procedure
Chinch Bug	Treat when bugs are causing stand reduction.				Check seedling rice, particularly fields bordering wheat.
Fall Armyworm, True Armyworm	Treat when 6 or more armyworms per square foot early season. Late season treat when fall armyworms are damaging flag leaf.				Early season watch rice bordering wheat for migration of true armyworms into field (damage can occur quickly when armyworms move in).
Grasshopper	Treat when damage is evident.				Watch field borders, particularly near grassy areas.
Greenbug	2 to 3 greenbugs per plant on 1- to 2-leaf stage rice.				General visual observation.
Stink Bug	5 stink bugs per 10 sweeps during first 2 weeks of panicle emergence. 10 stink bugs per 10 sweeps after first 2 weeks of panicle emergence.				Sample for stink bugs during the morning hours or late evening. Stink bug levels are difficult to estimate during high daytime temperatures.
Rice Water Weevil	Leaf Feeding Scar Method – Total Number Plants With Feeding Scars on New Leaves				Inspect the youngest leaf on 40 rice plants at each stop for adult feeding scars. Avoid areas with thin stand. Do not count older leaves with scars.
	Stop No.	Don't Treat – Stop Scouting When Total Is Less Than	No Decision – Keep Scouting When Total Is	Treat – Stop Scouting When Total Is More Than	
	1	No decision	Between	40	
	2	11	Between	56	
	3	28	Between	72	
	4	44	Between	89	
	5	61	Between	105	
	6	78	Between	122	
	7	94	Between	139	
8	111	Between	156		

RICE INSECT CONTROL

Insect	Insecticide	Formulation/Acre	Lb ai/Acre	Acres/Gallon	Application/Comments	Minimum Days From Last Application to Harvest
Chinch Bug	carbaryl Sevin 80 S Sevin XLR or 4 F	1 1/4-1 7/8 lb 2-3 pt	1.0-1.5	2.7-4.0	Treat when chinch bugs are causing stand reductions. DO NOT use Sevin prior to or within 15 days after propanil. Flooding prior to treatment will move chinch bugs up from the soil onto the plants and increase chances of control. Flooding alone may not control chinch bugs.	14
	γ -cyhalothrin (R) Prolex/Declare 1.25 CS Proaxis 0.5 CS	1.28-2.05 oz 3.2-5.12 oz	0.0125-0.02	62-100 25-40	Apply Karate Z, Prolex, Proaxis, Declare and Mustang Max in minimum of 5 gal of water per acre. DO NOT release flood water within 7 days after application of Karate Z, Prolex, Proaxis, Declare or Mustang Max.	21
	λ -cyhalothrin (R) Karate Z 2.08 CS (See generics, p. 26)	1.6-2.56 oz	0.025-0.04	50-80	Karate Z, Prolex, Proaxis, Declare and Mustang Max can be safely applied before or after herbicide applications.	21
	ζ -cypermethrin (R) Mustang Max 0.8 EC	2.64-4.0 oz	0.0165-0.025	32-48.5		14
Fall Armyworm, True Armyworm	γ -cyhalothrin (R) Prolex/Declare 1.25 CS Proaxis 0.5 EC	1.28-2.05 oz 3.2-5.12 oz	0.0125-0.02	62-100 25-40	Treat when six or more armyworms per square foot or two or more large larvae plus headcutting are found per square foot. If insecticide is applied before propanil, check herbicide recommendations for alternative control methods. Treatment will be most successful in late afternoon. Later in the season, treat when fall armyworms are present and damaging the flag leaf.	21
	λ -cyhalothrin (R) Karate Z 2.08 CS (See generics, p. 26)	1.6-2.56 oz	0.025-0.04	50-80		21
	methyl parathion (R) Methyl 4 EC	1-1.5 pt	0.5-0.75	5.3-8	Methyl parathion may be applied 14 days before or after propanil.	15
	ζ -cypermethrin (R) Mustang Max 0.8 EC	3.2-4.0 oz	0.02-0.025	32-40	Monitor rice fields adjacent to wheat for movement of true armyworms into seedling rice.	14
Short-Horned Grasshopper	Treat when damage is evident. Border treatments may be beneficial.					
	γ -cyhalothrin (R) Prolex, Declare 1.25 CS Proaxis 0.5 CS	1.28-2.05 oz 3.2-5.12 oz	0.0125-0.02	62-100 25-40	Karate Z, Prolex, Proaxis, Declare and Mustang Max can be safely applied before or after herbicide applications.	21
	λ -cyhalothrin (R) Karate Z 2.08 CS (See generics, p. 26)	1.6-2.56 oz	0.025-0.04	50-80		21
	ζ -cypermethrin (R) Mustang Max 0.8 EC	3.2-4.0 oz	0.02-0.025	32-40		14
	methyl parathion (R) Methyl 4 EC	1 pt	0.50	8	Methyl parathion may be applied 14 days before or after propanil.	15

RICE INSECT CONTROL

Insect	Insecticide	Formulation/Acre	Lb ai/Acre	Acres/Gallon	Application/Comments	Minimum Days From Last Application to Harvest	
Rice Stink Bug (Scout in morning for best results)	carbaryl Sevin 80 S Sevin XLR or 4 F	1 1/4-1 7/8 lb 2-3 pt	1.0-1.5	2.7-4.0	Check infestation levels weekly or bi-weekly following 75% panicle emergence using a 15-inch diameter sweep net. Apply insecticide when 2 stink bugs are found per square yard, when 5 or more stink bugs per 10 sweeps are present during the first 2 weeks after fields initially reach 75% panicle emergence or when 3 stink bugs are found per square yard or more than 10 stink bugs per 10 sweeps are present thereafter. Sampling stink bugs should be conducted between 8-10 a.m. and 6-8 p.m. to get the best estimate of the population. Repeat treatment as necessary to maintain control.	14	
	malathion Malathion 57% EC	1-1 1/2pt	0.625-0.94	5.3-8.0		7	
	methyl parathion (R) Methyl 4 EC	1-1.5 pt	0.5-0.75	5.3-8.0		Methyl parathion may be applied 14 days before or after propanil.	15
	γ-cyhalothrin (R) Prolex, Declare 1.25 CS Proaxis 0.5 CS	1.28-2.05 oz 3.2-5.12 oz	0.0125-0.2	62-100 25-40		Apply Karate Z in minimum of 5 gallons of water per acre. DO NOT release flood water within 7 days after application of Karate Z, Prolex, Proaxis, Declare or Mustang Max.	21
	λ-cyhalothrin (R) Karate Z 2.08 CS (See generics, p. 26)	1.6-2.56 oz	0.025-0.04	50-80			21
	ζ-cypermethrin (R) Mustang Max 0.8 EC	2.64-4.0 oz	0.0165-0.025	32-48.5			14
Rice Water Weevil (Adults)	γ-cyhalothrin (R) Prolex, Declare 1.25 CS Proaxis 0.5 CS	1.28-2.05 oz 3.2-5.12 oz	0.0125-0.02	62-100 25-40	Karate Z and Mustang Max can be safely applied before or after herbicide applications. Drill Seeded: Apply Karate Z, Prolex, Proaxis, Declare and Mustang Max within 10 days after permanent flood when adults are present. Water Seeded: Apply Karate Z, Prolex, Proaxis, Declare and Mustang Max within 7 days after permanent flood when adults are present; a second application may be necessary 5-7 days later. Karate Z, Prolex, Proaxis, Declare and Mustang Max prevent adults from laying eggs. Treat based on leaf scar treatment threshold using Karate for drill or water seeded.	21	
	λ-cyhalothrin (R) Karate Z 2.08 CS (See generics, p. 26)	1.6-2.56 oz	0.025-0.04	50-80		21	
	ζ-cypermethrin (R) Mustang Max 0.8 EC	3.2-4.0 oz	0.02-0.025	32-40		14	
Rice Water Weevil (Eggs)	diflubenzuron (R) Dimilin 2 L	12-16 fl oz	0.125-0.25	8.0-10.6	Drill Seeded: Apply Dimilin within 10 days after permanent flood when adults are present. Water Seeded: Apply Dimilin within 7 days after permanent flood.		

RICE INSECT CONTROL

Insect	Insecticide	Formulation/Acre	Lb ai/Acre	Acres/Gallon	Application/Comments	Minimum Days From Last Application to Harvest
Rice Water Weevil (Larvae)	chlorantraniliprole Dermacor X-100	1.5-6.0 oz/100 lb seed (see label)	0.06-0.08			
	thiamethoxam Cruiser 5 FS	0.03 mg ai/seed 3.3 oz/100 lb seed			DO NOT plant or sow Cruiser-treated seed by aerial application. Cruiser is NOT labeled for use in water-seeded rice. DO NOT use treated fields for aquaculture of edible fish or crustaceans. DO NOT exceed 120 lb seed per acre.	
Midge Larvae	There are currently no insecticides available for midge control in rice.				Midges can be a problem in water-seeded rice. To avoid midge problems, use presprouted seed, promptly apply seed as soon as field is flooded and avoid seeding during cool weather.	
Rice Stalk Borer	λ -cyhalothrin (R) Karate Z 2.08 CS	1.92-2.56 oz	0.03-0.04	50-67	Application at boot stage and/or at panicle emergence for suppression of white heads.	21
	γ -cyhalothrin (R) Prolex, Declare 1.25 CS Proaxis 0.5 CS	1.28-2.05 oz 3.2-5.12 oz	0.0125-0.02	62-100 25-40	Rice stalk borer is seldom a pest.	21
Aphids, Greenbug, Birdcherry-Oat Aphid	γ -cyhalothrin (R) Prolex, Declare 1.25 CS Proaxis 0.5 CS	1.28-2.05 oz 3.2-5.12 oz	0.0125-0.02	62-100 25-40	Karate Z, Prolex, Proaxis, Declare and Mustang Max may only provide suppression.	21
	λ -cyhalothrin (R) Karate Z 2.08 CS (See generics, p. 26)	1.6-2.56 oz	0.025-0.04	50-80	Experience has shown that insecticide applications are justified when two to three greenbugs per plant occur on rice in the one- to two-leaf stage.	21
	ξ -cypermethrin (R) Mustang Max 0.8 EC	3.2-4.0 oz	0.02-0.025	32-40		14
Grape Colaspis	thiamethoxam Cruiser 5 FS	0.03 mg ai/seed			DO NOT plant or sow Cruiser-treated seed by aerial application. Cruiser is NOT labeled for use in water-seeded rice. DO NOT use treated fields for aquaculture of edible fish or crustaceans. DO NOT exceed 120 lb seed per acre.	