

CANOLA DISEASES

Terry Kirkpatrick and Stephen Vann

Disease	Fungicide	Active Ingredient	FRAC Code*	Rate/Acre	Days to Harvest	Comments
Blackleg (<i>Leptosphaeria</i>)	Quadris	azoxystrobin	11	6 - 15.5 fl oz	30	Group 11 fungicide. See label.
Seedborne Blackleg Seedling Rhizoctonia Damping Off Seedling Blight (<i>Alternaria</i>)	Dynasty	azoxystrobin	11	0.1 - 3.75 fl oz/cwt (seed treatment)	See label.	Group 11 fungicide. See label.
Sclerotinia Stem Rot	Endura	boscalid	7	5 - 6 fl oz	21	Maximum of 2 applications per season. Group 7 fungicide. Apply at 20% - 50% bloom stage.
	Proline 480SC	prothioconazole	3	4.3 - 5.7 fl oz	36	Maximum of 2 applications per year. Apply at 20% - 50% bloom stage. Group 3 fungicide.
	Quadris	azoxystrobin	11	6 - 15.5 fl oz	30	Group 11 fungicide. See label.

*FRAC Code – Fungicides with the same FRAC Code have the same mode of action. See <http://www.frac.info/frac/index.htm> for an explanation of the FRAC Codes. Rotation of fungicides with different FRAC Codes could minimize the development of fungicide-resistant strains.