

PEANUT SEED TREATMENTS (Commercial Production)

Cliff Coker and Terry Kirkpatrick

| Disease | Fungicide | Active Ingredient | FRAC Code* | Rate/cwt Seed | Comments |
|---|-----------------------------|-------------------|----------------|------------------|---|
| <i>Rhizoctonia solani</i> <i>Pythium</i> | Dynasty PD | azoxystrobin + | 11 | 3 - 4 oz | For use by commercial seed treaters only. |
| | | mefenoxam + | 4 | | |
| | | fludioxonil | 12 | | |
| | Maxim 4FS | fludioxonil | 12 | 0.08 oz | |
| | Trilex Optimum | trifloxystrobin + | 11 | --- | |
| metalaxyl + carboxin | | 4 | | | |
| Trilex Star | trifloxystrobin + | 11 | --- | | |
| | metalaxyl + | 4 | | | |
| | carboxin | --- | | | |
| Uniform | azoxystrobin + mefenoxam | 11 4 | 0.34 oz/row ft | In-furrow spray. | |
| <i>Pythium</i> | Apron XL | mefenoxam | 4 | 0.32 oz | |

*FRAC Code – Fungicides with the same FRAC Code have the same mode of action. See <http://www.frac.info/frac/index.htm> for an explanation of the FRAC Codes. Rotation of fungicides with different FRAC Codes could minimize the development of fungicide-resistant strains.