

SMALL FRUIT DISEASES (Home Garden)

Stephen Vann

Disease	Product	Active Ingredient	Rate	Comments
Fungal leaf and fruit diseases on grapes, blackberry, raspberry and strawberry	Ortho Home Orchard Spray	captan	5 tbs/gal	Spray on a 7 - 14 day schedule from early bloom until harvest, depending on weather and disease. See labels for additional information. Good coverage is essential.
	Hi-Yield Captan 50WP	captan	2 tbs/gal	
	Hi-Yield Bordeaux Fungicide Mix	copper salts	See label.	Apply at first sign of disease, 10 - 14 day interval. See label.
	Bayer Advanced Natria Disease Control	<i>Bacillus subtilis</i>	RTU*	Spray to runoff. See label.
	Monterey Liqui-Cop	copper	See label.	See label.
	Bonide Copper Liquid Fungicide Concentrate	copper salts	2 - 4 tsp/gal	See label.
	Bonide Lime Sulfur Spray	calcium polysulfide	See label.	See label.
	Bonide Sulfur Plant Fungicide	sulfur	1 - 3 tbs/gal	See label.
	Safer Brand Concern Garden Defense Conc.	neem oil extract	1 fl oz/gal	7 - 14 days. Complete coverage important. RTU* also available. Organic uses.
	Bonide Citrus, Fruit and Nut Orchard Spray Concentrate	sulfur + pyrethrins	2.5 - 5 oz/gal	7 - 10 days. Apply at first evidence of disease.
Natural Guard Copper Soap Liquid Fungicide	copper octanoate	0.5 - 2 oz/gal	7 - 10 days. Apply at first evidence of disease.	
Fungal diseases of <u>grapes only</u>	Spectracide Immunox Plus	myclobutanil + insecticide	See label.	Spray on a 14 day schedule from very early bloom until 2 weeks before harvest. See label.
	Hi-Yield Bordeaux Fungicide Mix	copper salts	8 - 9 tbs/gal	Apply in early spring. See label precautions.
	Bonide Mancozeb Flowable with Zinc	mancozeb	3 - 4 tsp/gal	Begin at 1/2 inch new growth, continue at 7 - 10 days until bloom. See label.

*RTU = Ready To Use