

Measuring Forage Moisture Content Using an Air Fryer

John Jennings
Professor - Forages
Animal Science

Measuring moisture content of forage cut for hay or silage is an essential step to ensure storage stability and product quality. Hay baled with too much moisture can mold or be subject to spontaneous heating. Silage baled or chopped at moisture contents outside a recommended range may not ferment properly, reducing storage life and animal acceptance. A relatively new method of measuring forage moisture content is through use of an air fryer. This household appliance is basically a small convection oven. It can be used at the farm shop or can be operated in the field from a generator to provide accurate forage moisture readings.

4. Set the air fryer to 250°F for 30 minutes.
5. Weigh the dried sample and calculate the dry matter content.

Calculating Hay Moisture:

Remember that for 100-gram samples, the number of grams of moisture lost equals the percent moisture.

Example: A 100-gram sample was dried and the final weight was 80 grams.

$$100 - 80 = 20 \text{ grams moisture lost} \\ = 20\% \text{ moisture}$$

If other than a 100-gram sample is used, use this equation to calculate hay moisture content:

$$\frac{\text{Starting weight (grams)} - \text{final weight (grams)}}{\text{starting weight (grams)}} \times 100 = \% \text{ Moisture}$$

Example: The starting weight was 90 grams and the final weight of the sample is 70 grams. Subtract 70 from 90 grams (starting weight) to get the amount of moisture lost during drying. Divide the amount of moisture lost by the starting weight and multiply by 100 to get the final sample moisture percent.

$$90 - 70 = 20 \text{ grams moisture lost} \\ \frac{20}{90} \times 100 = 22\% \text{ moisture}$$

Steps for Using an Air Fryer to Measure Hay Moisture

Materials Needed:

- air fryer
- gram scale
- wire screen to hold sample in place during drying (needed on some models)
- plate or bowl to contain sample for weighing
- calculator

1. Take a core sample from test bales or cut up a representative hay sample into small pieces (less than 1 inch) and weigh 100 grams onto a paper plate.
2. Pour the sample into the air fryer.
3. Place a screen over the sample to hold it in place during drying.

*Arkansas Is
Our Campus*

Visit our website at
<https://www.uaex.uada.edu>

Printed by University of Arkansas Cooperative Extension Service Printing Services.

DR. JOHN JENNINGS is professor - forages with the Department of Animal Science, University of Arkansas System Division of Agriculture, Little Rock.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director, Cooperative Extension Service, University of Arkansas. The University of Arkansas System Division of Agriculture offers all its Extension and Research programs and services without regard to race, color, sex, gender identity, sexual orientation, national origin, religion, age, disability, marital or veteran status, genetic information, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.

FSA3155-PD-10-20N