

Forage, Feed and Grazing Restrictions for Soybean Herbicides

Herbicide	Restrictions
Assure II	Do not graze treated fields or harvest for forage or hay. Do not apply within 80 days of harvest.
Authority MTZ	Do not graze or harvest for forage or hay.
Basagran/Broadloom	No restrictions on label.
Broadstrike + Dual or Treflan, Python	Do not graze or feed treated soybean forage, hay or straw to livestock.
Butyrac/Butoxone (2,4-DB)	Do not graze or feed soybean hay within 60 days after application of a 2,4-DB tank-mix application. Do not harvest soybeans within 60 days after spray application.
Canopy	Do not graze treated fields or harvest for forage or hay.
Classic	Do not graze treated fields or harvest for forage or hay. Do not apply later than 60 days before soybean maturity.
Cobra	Do not graze animals on green forage or stubble. Do not utilize hay or straw for animal feed or bedding. Apply not later than 90 days before harvest.
Dual Magnum	No restrictions on label.
Enlist One	Do not graze or harvest for forage or hay.
Enlist Duo	Do not graze or harvest for forage or hay.
Engenia	Do not feed soybean fodder or hay following a preharvest application.
FirstRate	Do not harvest for forage or hay for 25 days.
Fusilade DX, Fusion	Do not graze or harvest for forage or hay.
Glufosinate	Do not feed green soybean plants to livestock.
Glyphosate	Do not harvest or feed treated crops for 8 weeks after application. Allow 14 days following spot treatment or selective equipment before grazing domestic livestock.
Metribuzin	Treated vines* may be grazed or fed to livestock 40 days after application.
Outlook	Do not graze or feed forage, hay or straw to livestock.

Herbicide	Restrictions
Paraquat	Do not graze treated areas or feed treated forage to livestock.
Poast/ Poast Plus	Do not graze treated fields and do not feed treated soybean forage (green succulent) or ensilage to livestock. Treated soybean hay may be fed. Do not apply to soybeans within 90 days of harvest.
Prowl	Livestock <i>can</i> graze or be fed soybean forage from treated fields.
Pursuit	Do not graze or feed treated soybean forage, hay or straw to livestock. There should be an interval of at least 85 days between an application of Pursuit and soybean harvest.
Python	Do not graze or harvest for forage or hay.
Reflex/Flexstar/ Sinister	Do not graze treated areas or harvest for forage or hay. Do not graze rotated small grain crops or harvest for livestock forage or straw. Do not apply after first bloom.
Resource	Do not graze or harvest for forage or hay.
Select/TapOut	Do not graze treated fields or feed treated forage or hay to livestock.
Storm	Do not use treated plants for feed or forage. Do not apply within 50 days of harvest.
Surveil	Do not graze or harvest for forage or hay.
Synchrony	Do not graze treated fields or harvest for forage or hay. Do not apply later than 60 days before soybean maturity.
Tavium	Do not graze or feed to livestock, or harvest for food, any cover crop planted following an application.
Treflan	No restrictions on label.
Trivence	Do not graze or harvest for forage or hay.
Typhoon	Do not graze or harvest for forage or hay.
Ultra Blazer	Do not use treated plants for feed or forage.
Valor	Do not graze or harvest for forage or hay.
Warrant	Do not graze or feed for 60 days.
Xtendimax	Do not feed soybean fodder or hay following a preharvest application.
Zidua/Anthem	Do not graze or feed for 37 days.

Restrictions are listed as worded on the labels. Feeding and application restrictions for herbicides are generally based on residue tolerances allowed for animal feeding. The restrictions are generally not due to acute toxicity (poisoning) problems. Livestock that are accidentally fed treated crops earlier than allowed will not be harmed, but may have illegal pesticide residues in their meat or milk. If you have fed livestock treated crops within the restricted period, refer to the label, your dealer or herbicide company representative for more information.

* Many labels refer to soybean plants as 'vines'.