

Footnotes (continued)

- ¹¹Increase rotational interval for Grain Sorghum to 18 months if pH is 7.5 or greater or if Capreno rate exceeds 3 oz/A.
- ¹²Time interval is based on 8 oz/A application rate and does not begin until 1 inch of rainfall is received.
- ¹³Clearfield Rice may be planted at any time after application of Beyond, Clearpath or Newpath.
- ¹⁴If 4 oz/A or less used and 1 inch of rainfall/irrigation received after application.
- ¹⁵Conduct field bioassay prior to planting these crops.
- ¹⁶STS Soybeans can be planted immediately.
- ¹⁷As fallow bed treatment. Prior to planting, treated fallow beds should be thoroughly tilled (incorporated) to a depth of at least 2.5 inches.
- ¹⁸Wheat can be planted the following year after application of Harness Extra.
- ¹⁹Rotation interval for Soybeans is 2 months where pH is less than 7.5.
- ²⁰If less than 15 inches of rainfall received since application, extend replant intervals to 18 months. If pH greater than 6.5, do not plant Rice following year.
- ²¹Rotation intervals increase with higher application rates.
- ²²18 months for Cotton if rate is greater than 5 oz/A and pH >7.2.
- ²³Rotation to Grain Sorghum is 18 months when Spartan is applied at 8 oz/A.
- ²⁴Rotation to Cotton and Grain Sorghum is 18 months when Spartan Charge is applied at 10.2 fl oz/A.
- ²⁵10 months for 1 oz/A, 12 months for 2 oz/A, 18 months for 3 oz/A and 24 months for 4 oz/A.
- ²⁶4 month rotation to Soybeans if 4 oz/A rate is used.
- ²⁷Enlist Corn, Cotton and Soybeans can be planted immediately.
- ²⁸Xtend Soybean and Cotton can be planted immediately.
- ²⁹Most 2,4-D labels state rotation to all crops after 90 days or sufficiently dissipated.
- ³⁰Days listed are based on University data and after receiving 1 inch of rainfall.

Authors

Tom Barber, Professor - Weed Science
Jason Norsworthy, Professor - Weed Science

University of Arkansas System Division of Agriculture
Weed Science Program



2018
Row Crop Plant-Back
Intervals for
Common Herbicides



DR. TOM BARBER with the Crop, Soil and Environmental Sciences Department, University of Arkansas System Division of Agriculture, is located at Lonoke. **DR. JASON NORSWORTHY** also with the Crop, Soil and Environmental Sciences Department is located at the University of Arkansas, Fayetteville.

University of Arkansas, United States Department of Agriculture,
and County Governments Cooperating

Printed by University of Arkansas Cooperative Extension Service Printing Services.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director, Cooperative Extension Service, University of Arkansas. The University of Arkansas System Division of Agriculture offers all its Extension and Research programs and services without regard to race, color, sex, gender identity, sexual orientation, national origin, religion, age, disability, marital or veteran status, genetic information, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.

MP519-PD-3-2018RV



The following table consists of herbicide trade and common names and their labeled plant-back restrictions for six major row crops grown in Arkansas. Designations for time periods are as follows:

I – crops can be planted immediately, D – the number of days for crop replant interval, M – the number of months, FY – crops can be planted the following year and FS – the following season. Brand names are subject to change; always read and follow label instructions.

Herbicide	Cotton	Field Corn	Grain Sorghum	Rice	Soybeans	Wheat
Accent/Zest	10M	I	10-18M ¹	10-18M ²	15D	4M
Aim	I	I	I	I	I	I
Anthem	4M	I	11M	10M ³	I	4M
Anthem ATZ	FY	I	FY	FY	FY	FY
Anthem Flex	I	I	11M	10M	I	I
Armezon	9M	I	9M	3M	9M	3M
Assure II	I	120D	120D	120D	I	120D
Atrazine ⁴	FY	I	I	FY	FY ⁵	FY
Authority Elite	12-18M ⁶	10M	10M	10M	I	4.5M
Authority MTZ/XL	12M ⁶	10M ⁶	10M ⁶	10M ⁶	I	4M
Axial XL	90D	90D	90D	90D	90D	I
Axiom	8M	I	1Y	12M	I	7D
Balance Flexx	10M	I	6M	10M	6M	4M
Basagran	I	I	I	I	I	I
Beyond	9M	8.5M	9M	9M ¹³	I	3M
Bicep II/Magnum ⁴	FY	I	I ⁸	FS	FY	15M
Bolero	6M	6M	6M	6M	6M	6M
Boundary	12M	8M	12M	8M	I	4.5M
Cadre	18M	9M	18M	26M	9M	4M
Callisto	10M	I	I	FY	10M	4M
Canopy DF	10M ⁶	9M ⁶	10M ⁶	10M ⁶	I	4M
Caparol 4L	I	FY	FY	FY	FY	FY
Capreno ¹¹	10M ⁹	I	10M ⁹	10M ¹⁰	10M ⁹	4M
Clarity	21D ¹²	I	15D	22D	14D ¹²	22D
Classic	8M	8M	9M	9M	I	3M
Clearpath ⁷	18M	10M	18M	18M ¹³	10M	10M
Clethodim/Select	I	30D	30D	30D	I	30D
Clincher SF	3M	3M	3M	I	3M	3M
Cobra	I	I	I	I	I	I
Command	I	9M	9M	I	I	4M ²¹
Corvus	10M ⁹	I	17M ¹⁰	10M ⁹	9M ⁹	4M
Cotoran	I	6M	6M	6M	6M	6M
Direx	I	I	6M	6M	6M	6M
Distinct	30 ¹⁴	7D	30 ¹⁴	30 ¹⁴	30 ¹⁴	30 ¹⁴
Dual/Magnum	I	I	I ⁸	FS	I	4.5M
Duet	60D	60D	60D	I	60D	60D
Elevore	30D	14D	14D	14D	14D	14D
Engenia ²⁸	21-42D	I	14-28D ^{21,30}	120D	I	120D
Enlist One	30D ²⁷	14D ²⁷	7D ³⁰	21D	15-30D ^{21,27}	7D
Enlist Duo	30D ²⁷	7-14D ^{21,27}	7D ³⁰	21D	30D ²⁷	7D
Envive ⁶	10M	10M	10M	9M	I	4M
Envoke	7M	7M	7M	7M	7M	3M
Facet L	10M	10M	I	I	10M	I

Herbicide	Cotton	Field Corn	Grain Sorghum	Rice	Soybeans	Wheat
FeXapan ²⁸	22-120D ^{21,28}	22-120D ²¹	22-120D ²¹	22-120D ²¹	22-120D ^{21,28}	22-120D ²¹
Fierce	45D	30D	12M	10M	I	1-2M
Finesse G&B	FB ¹⁵	14M	FB	FB	6 ¹⁶	I
FirstRate	9M	9M	9M	9M	I	4M
FirstShot	14D ⁶	14D ⁶	14D ⁶	I	1D ⁶	I
Flexstar	I	10M	10M	10M	I	4M
Fusilade DX/Fusion	I	60D	60D	60D	I	60D
Gambit	10M	I	2M	I	10M	2M
Gangster/Surveil	9M	9M	9M	9M	I	3M
Glyphosate	I	I	I	I	I	I
Goal 2XL	7D ¹⁷	30D	10M	10M	7D ¹⁷	10M
Gramoxone SL	I	I	I	I	I	I
Grandstand	4M	4M	4M	I	4M	4M
Grasp/Xtra	3M	3M	3M	3M	3M	3M
Guardman Max	FY	I	I	2Y	FY ⁵	1Y
Halex GT	10M	I	I	FY	10M	4M
Harmony Extra	14D ⁶	14D ⁶	14D ⁶	I	7D ⁶	I
Harmony GT	7D	I	I	I	I	I
Harness	FY	I	FY	18M	FY	4M ¹⁸
Hoelon	FS	FS	FS	FS	FS	I
Huskie	FY	9M	4M	FY	4M	1M
Keystone NXT	FY	I	I	FY	FY	4M
Laudis	10M	I	10M	10M	8M	4M
Layby Pro	4M	4M	4M	4M	4M	3M
Leadoff	30D ⁶	I	10M	10M	30D ⁶	3M
League	8M	12M	12M	I	12M	12M
Lexar EZ	FS	I	FS	FS	FS	FS
Liberty	I	I	180D	I	I	70D
Linex	4M	I	I	1Y	I	4M
Londax	4M	4M	4M	I	4M	4M
Loyant	3M	3M	3M	I	3M	3M
Marksman ⁴	FY	I	I	FY	FY ⁵	10M
MSMA	I	I	I	I	I	I
Newpath ⁷	18M	8.5M	18M	18M ¹³	I	4M
Obey	309D	309D	309D	I	309D	309D
Osprey	90D	12M	10M	90D	90D	7D
Outlook	FY	I	I ⁸	FY	I	4M
Paraquat	I	I	I	I	I	I
Peak	10M	1M	I	10M	10M	I
Permit/HaloMax	4M	1M	2M	I	9M ^{16,19}	2M
Permit Plus	4M	1M	2M	I	9M ^{16,19}	2M
Poast Plus	I	30D	30D	30D	I	30D
PowerFlex HL	90D	9M	90D	1Y	90D	1M
Prefix	1M	10M	10M	10M	I	4.5M
Provisia	I	120D	120D	120D	I	120D
Prowl H ₂ O	I	FS	FS	FS	I	4M
Python	18M	I	12M	6M	I	4M

Herbicide	Cotton	Field Corn	Grain Sorghum	Rice	Soybeans	Wheat
Realm Q	10M	I	10M	18M	10M	4M
RebelEX	30D	30D	30D	I	30D	30D
Reflex	I	10M	10M	10M	I	4M
Regiment	FY	FY	FY	I	FY	FY
Resolve Q	30D	I	10M ²⁰	10M ²⁰	30D	3M
RiceBeaux	60D	60D	60D	I	60D	60D
RicePro	309D	309D	309D	I	309D	309D
Rice-Star HT	30D	30D	30D	I	30D	4M
Select Max	I	30D	30D	30D	I	30D
Sencor	8M	4M	18M	8M	I	4M
Sequence	I	I	I	FS	I	4M
Sharpen	1.5-3M ²¹	I	I	I	0-30D ²¹	I
Solicam	30D	2Y	2Y	2Y	45D	2Y
Sonic/Authority First	12-18M ²²	10M	12M	10M	I	4M
Spartan	18M	10M	10M ²³	10M	I	4M
Spartan Charge	12M ²⁴	4M	10M ²⁴	10M	I	4M
Stam M4	I	I	I	I	I	I
Staple LX	I	10M	18M	9M	10M	4M
Storm	100D	100D	100D	I	I	40D
Strada PRO	6M	3M	3M	I	9M	3M
Strada XT ⁷	309D	309D	309D	I	309D	309D
Suprend	7M	7M	7M	7M	7M	3M
Surestart II	26M	I	12M	FY	FY	4M
Surpass	2Y	I	FS	2Y	FS	4M
Treflan	I	12M	12M	12M	I	12M
Ultra Blazer	100D	100D	100D	I	I	40D
Valor	30D ⁶	30D ⁶	30D ⁶	30D ⁶	I	30D ⁶
Verdict	1.5-4M ²¹	I	I ⁸	FY	0-4M ²¹	4M
Warrant	I	I	I	FY	I	4M
Xtendimax ²⁸	21D	14-28D ²¹	15D	120-180D ²¹	14-28D ²¹	22-45D ²¹
Zidua	0-4M ²¹	I	6-12M ²¹	10-24M ²⁵	I ²⁶	0-6M ²¹
2,4-D ²⁹	28D	7D ³⁰	7D	21D	14D	7D

¹ Grain Sorghum may be planted 10 months after application on soils with pH 7.5 or less. If soil pH is greater than 7.5, do not plant Grain Sorghum less than 18 months after application.
² Rice may be planted 10 months after Accent application on soils with pH 6.5 or less. If soil pH is greater than 6.5, do not plant Rice less than 18 months after application.
³ 3.25 oz use rate, 12 months for 6.5 oz/A, 18 months for 9.75 oz/A.
⁴ If applied after June 10, do not rotate to any crop other than Corn or Grain Sorghum the year following application or injury may occur.
⁵ Injury may occur to Soybeans planted in soils with a calcareous surface layer.
⁶ Time interval increases with the increase of rate and soil pH level.
⁷ For Newpath use rates greater than 8 oz/A per season; only Soybeans may be planted the following year.
⁸ Replant only with Concept-treated or screen-treated seed.
⁹ Needs 15 inches cumulative precipitation from application to planting rotational crop.
¹⁰ Needs 30 inches cumulative precipitation from application to planting rotational crop.

(footnotes continued on next panel)