

### Crop Replant and Rotation Guide for Rice Herbicides

Herbicide	Replant/Crop Rotation	Time Interval	Precautions
2,4-D	All	90 days	90 days or until sufficiently dissipated.
Aim	AL, CL All other	12 months I	
Basagran	All	I	
Beyond	S A B, CT, GS, O, SF, P, C, R CA	I 3 months 9 months 18 months	
Bolero	All*		* Do not plant subsequent crops in treated fields within 6 months of last application. Do not use in fields where fall farming of crayfish will be practiced. Do not apply to second stubble rice crop.
Clincher	All	3 months	No more than 25 oz per growing season. Keep away from peach trees.
Command	S W	I 4 months	
Facet	R All	I 309 days	Do not plant tomatoes or carrots within 2 years. Do not use in fields for fish farming or where fall farming of crayfish will be practiced.
Grandstand	All	6 months	
Grasp	All	3 months	
League	R CT C, S, GS	I 8 months 12 months	* See label for vegetables and other crops.
Londax	All	120 days	
Loyant	All	3 months	
Newpath	S, P, Clearfield Corn, Clearfield Rice W C CT, GS, Non-Clearfield Rice	I 4 months 8.5 months 18 months	Wheat rotation is longer if more than 8 oz/year is used.
Obey	R All	1 309 days	Do not plant tomatoes or carrots for 2 years. Do not use in fields where fish or crayfish will be farmed.

Herbicide	Replant/Crop Rotation	Time Interval	Precautions
Permit, Permit Plus or Halomax	R W, C, GS CT, P S SF	I 2 months 6 months 9 months 18 months	Preplant to rice up to pH 8.
Propanil	All	I	
Prowl	CT, S All B, W	I FY 120 days*	* 90 days after a post incorporated application in irrigated field corn or grain sorghum. Do not plant following irrigation if crop failure occurs. Do not plant following no-till practices. Do not plant following higher Prowl rates for rhizome johnsongrass or red rice.
Regiment	R All	I FY	
Ricestar HT	C, CT, GS, S SG	30 days 120 days	
Ultra Blazer	P, S All*	I FY	* Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in treated fields for a period of 18 months following treatment.
Sharpen	C, R, GS, SG, S CT All	I 1.5 months 4 months	Sweet corn plant back is 0.5 months.
Storm	All	Fall	
Strada	R C, SG C, S All others	I 3 months 6 months 12 months	

\* This table applies to the major field and forage crops. Refer to the herbicide labels for the latest recrop and rotation information for horticultural crops. These are written as best we could interpret the labels. We regret any omissions or errors. Always refer to product labels before using a pesticide or replanting into treated fields.

#### Key Crop

All = All crops not specified	GS = Grain Sorghum	S = Soybeans
B = Barley	O = Oat	SF = Sunflowers
C = Corn	P = Peanuts	SG = Small Grains
CA = Canola	R = Rice	W = Wheat
CT = Cotton		

#### Timing

I = Immediately                      FY = Following year (usually spring)