

**WEED RESPONSE RATINGS FOR RICE HERBICIDES (cont.)**  
(See Explanation for Ratings Tables on Page 3.)

HERBICIDES	HERBICIDE FAMILY	GRASSES								BROADLEAF WEEDS														SEDGES					
		Barnyardgrass <sup>1</sup>	Broadleaf Signalgrass	Crabgrass	Fall Panicum	Red Rice	Rice Cutgrass	Sprangletop (loosehead) (bearded sprangletop)	Sprangletop (tighthead) (Amazon)	Ammania (red stem)	Dayflower	Ducksalad	Eclipta	False Pimpernel	Gooseweed	Groundcherry	Hemp Sesbania (coffeebean)	Indian Jointvetch	Northern Jointvetch (curly indigo)	Palmleaf Morningglory	Pigweed, Palmer	Pitted Morningglory	Smartweed	Texasweed	Water Hyssop	Flatsedges	Spikerush	Umbrella Sedge	Yellow Nutseed
<b>Midseason</b>																													
2,4-D	4	0	0	0	0	0	0	0	0	9	9	9	9	9	6	5	9	5	5	9	8	9	6	0	9	8	8	3	5
2,4-D + propanil for levees	4,7	6	6	2	6	0	0	6	6	9	9	8	9	9	8	5	9	8	8	8	9	9	7	0	9	8	8	3	6
Grandstand + propanil	4,7	4	4	4	4	0	0	0	0	9	-	6	6	8	7	3	9	8	9	9	7	9	5	0	8	5	8	5	3
Propanil	7	4	4	4	4	0	0	0	0	4	0	3	4	4	0	4	8	5	5	3	6	0	3	0	8	5	7	5	3
Propanil + Ultra Blazer	7,14	5	5	5	5	0	0	0	0	5	2	4	5	5	2	5	9	6	6	7	7	8	7	0	8	6	7	5	4
Ultra Blazer	14	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	9	0	0	3	6	3	6	0	0	0	0	0	0

\*Water seed pin-point flood culture.    \*\*Postemergence control only.    <sup>1</sup>Some biotypes of barnyardgrass in Arkansas are resistant to Command, propanil, Facet or both (Facet + propanil), and Newpath, Grasp, and Regiment.  
Rating Scale – 0 = No Control    10 = 100% Control.

**Rainfall-free Periods for Postemergence Rice**

Herbicide	Time Before Rainfall	Herbicide	Time Before Rainfall
2,4-D	6 hrs	Newpath	4 hrs
Aim	1 hr	Permit/Permit Plus	4 hrs
Basagran	8 hrs	Propanil	6 hrs
Beyond	4 hrs	Regiment	8 hrs
Bolero	nothing on label	Ricestar HT	1 hr
Clincher	1 hr	Sharpen	1 hr
Facet	nothing on label	Storm	8 hrs
Grasp	1 hr	Strada	6 hrs
League	6 hrs	Ultra Blazer	4 hrs