



This bulletin from the Cooperative Extension Plant Health Clinic (Plant Disease Clinic) is an electronic update about diseases and other problems observed in our lab each month. Input from everybody interested in plants is welcome and appreciated.

Cottonwood

The Eastern Cottonwood is one of the largest North American hardwood trees, growing to 20–40 m tall and with a trunk up to 1.8 m diameter. They are closely related to Aspens and Poplars, sharing many of the same diseases. Leaf and shoot blight caused by *Venturia* spp. rarely seriously harms healthy, established trees, but can cause serious damage to saplings. Severe infections can reduce growth. After trees reach a height of 15 feet, damage is minimal. Symptoms are brown to black irregular lesions on leaves and stems early in the spring. Damaged stems become black, brittle, and curl to resemble a shepherd's crook. Recently infected tissues will have an olive green look from spore masses. Fallen leaves should be raked up and disposed of away from the property. Severely affected trees can be sprayed with a fungicide containing Thiophanate-methyl, such as Cleary's 3336. In most cases, chemical controls are not warranted.

Strawberry by Rebecca Barocco

Strawberry leaf scorch is a disease limited to *Fragaria* species. The leaves are the primary source of infection displaying dark purple circular spots that can develop a reddish brown center tinge. These lesions will coalesce and become scorched by turning brown and drying the tips so much that they begin to curl. Symptoms may also occur on leaf stems as well as flower and fruit parts with somewhat similar lesions to those of the leaves. The enemy here is the ascomycete fungus *Diplocarpon earlianum*. The ascospores are dispersed by air which may come from another strawberry source, possibly from the wild, or from dead plant residue that overwinters the fungus in black structures called apothecia. These apothecia can be seen as raised black spots on the leaf surface. Because of this overwintering mechanism, fallen dead tissue should be removed from the ground to prevent further infection. Spores that land on the tissue germinate and grow under wet conditions during temperatures from 36-86°F with

Cottonwood leaf blight



Sherrie Smith University of Arkansas Cooperative Extension

59-77°F being optimal. The conidiospores cause the continual infection during the season and are spread by splashing water. There are resistant cultivars that can be grown, but if these are not desirable, then other control measures can be taken. Generally the first and second years of planting are not as susceptible, so new plantings can help prevent the disease. Other methods are holding off on nitrogen application in the spring and only applying optimal levels during summer renovation, planting on a light slope for air drainage, and not overcrowding within rows. Planting in a sunny area with irrigation practices that prevent long periods of leaf wetness, such as not using overhead irrigation or watering in the morning, will inhibit fungal growth. If economically feasible, a fungicide should be sprayed first in the spring between bud break and bloom with either Captan or Thiram and then again every 7 to 10 days until 14 days before harvest with either Elevate, Captan, Thiram, or Switch. For the rest of the season,



spray every 2 to 3 weeks with either Captan, Thiram, Topsin, or Syllit. Homeowners can use products with captan or with the organic fungicides containing copper, calcium polysulfide, sulfur, or sulfur and pyrethrins. However, copper can cause burning of the tissue. Labels should be read carefully for organic substances to prevent such damage.

Strawberry scorch



Sherrie Smith University of Arkansas Cooperative Extension

Mulberry

Mulberry trees are grown around the globe for their fruit, lumber, and for their value in the landscape. Mulberry trees are prone this time of year to a fungal leaf spot disease caused by *Cercospora mori* or *Cercospora mori*. The initial symptoms of this disease are small dark spots in early spring that gradually increase in size through the growing season. The spots gradually become circular with the centre appearing as a white dot somewhat like a halo, the margins remain dark brown, with spores developing in the lesions. Severe infections cause defoliation which can weaken a tree already under stress. Weeping mulberries are small enough to be easily sprayed. Clean up all fallen leaves and spray with an ornamental fungicide or treat with Bayer Systemic

Fungicide. Those growing mulberries for the fruit can use an orchard spray. Begin at green tip in the spring.

Mulberry Cercospora



Brad McGinley University of Arkansas Cooperative Extension



Sherrie Smith University of Arkansas Cooperative Extension

Mulberry fruit also can get a disease called Popcorn disease, caused by the fungus, *Ciboria carunculoides*. The symptoms appear only on infected fruit. Individual Carpels of the fruit are replaced by sclerotia, which enlarge and extend beyond healthy berries. White mulberry varieties and hybrids are more susceptible to popcorn disease.



This disease is not considered economically important on ornamental mulberries. It can be a problem, however, on mulberries propagated for fruit production and can cause high yield losses. Control is achieved by taking sanitary measures. Remove and bury the infected fruit on the trees and any ground debris as it appears during the growing season.

Popcorn disease



Steve Vann University of Arkansas Cooperative Extension

APLPV



Sherrie Smith University of Arkansas Cooperative Extension

APLPV



Sherrie Smith University of Arkansas Cooperative Extension

Plum

A plum sample tested positive for American Plum Line Pattern Virus (APLPV) per an AGDIA crop screen. APLPV causes ringspots and line patterns on plum and peach cultivars. The vector for transmission is unknown, but appears easily transmissible by bud grafting and mechanical means. Symptoms are diffused lines, bands, oak-leaf patterns, or rings. The literature states that symptoms are confined to the leaves. The sample tested showed striking fruit symptoms as well as leaf ringspots. However, very similar symptoms are induced by Apple Mosaic Virus and certain strains of Prunus Necrotic Ringspot Virus, neither of which is serologically related to American plum line pattern virus. The virus has a wide host range; it is known to infect 85 of 245 species inoculated with sap, including six woody rosaceous species. Virus is not curable.