



This bulletin from the Cooperative Extension Plant Health Clinic (Plant Disease Clinic) is an electronic update about diseases and other problems observed in our lab each month. Input from everybody interested in plants is welcome and appreciated.

## Corn

Weather conditions this growing season have been ideal for corn diseases. **Diplodia ear rot** is most severe when corn follows corn and wet weather occurs shortly after silking. The disease is caused by the fungus, *Stenocarpella maydis*. Symptoms are bleached to straw colored husks. A conspicuous gray to white mycelial growth may be observed over the entire ear. The growth typically starts at the base of the ear and moves upward. The ear may appear shrunken with the infected kernels glued to the husk by the fungal growth. Late in the season, black pycnidia may be observed on the husks, kernels, cobs, and rotted stalks. *Stenocarpella maydis* is also responsible for an important stalk rot of corn. It can be distinguished from other stalk rots by the presence of sub epidermal, minute, dark brown to black pycnidia in the rind tissue of the lower stem. The two most important methods of reducing the incidence of Diplodia ear and stalk rot are crop rotation and fall tillage of corn residue.

### Diplodia stalk rot



D.G. White, APS Image Library

### Diplodia ear rot



Sherrie Smith University of Arkansas Cooperative Extension

**Gibberella stalk rot**, caused by *Gibberella maydis*, has symptoms that are similar to those of other stalk rots. Plants wilt, the leaves change to a dull green, and lower stalks become straw colored. Red discoloration inside the stalk and disintegration of the pith are diagnostic for *Gibberella* stalk rot.



## Gibberella stalk rot



Sherrie Smith University of Arkansas Cooperative Extension

**Charcoal rot** symptoms are similar to other stalk rots. It can be differentiated by the presence of numerous, minute, black sclerotia on the vascular bundles and inside the rind, causing the interior of the stalk to appear gray-black. The causal organism is *Macrophomina phaseolina*, the same fungus that causes charcoal rot in beans. Root stress related to drought and/or too much water can cause charcoal rot.

**Anthrax stalk rot**, caused by *Colletotrichum graminicola*, is recognized late in the season by the shiny black color on the outer stalk. The black color may be uniform or blotchy. The stalk can be easily crushed at the point of discoloration. The pathogen may rot several internodes on the stalk. Balanced fertility and good water management reduces the incidence of

anthracnose when coupled with cultivars with some resistance.

## Charcoal rot



D.G. White, APS Image Library

## Charcoal rot sclerotia



D.G. White, APS Image Library




## Anthracnose stalk rot



D.G. White, APS Image Library

## Corn red leaf



Rebecca Barocco University of Arkansas Cooperative Extension  
Plant Health Clinic Intern



## Corn-abiotic

Red leaves and/or red stems with barren stalks may be caused by a broken midrib, or low fertility, dense plant populations, aphid or herbicide damage, chewed off silks, or poor timing of silking and pollen shed. An accumulation of sugars and other photosynthetic products in leaves and sheaths of barren stalks produces the red coloration.

## Northern corn leaf blight


Northern corn leaf blight is common here in the Mid-south with our warm, humid summers. The disease, is caused by *Setosphaeria turcica*, and causes gray-green, elliptical or cigar-shaped lesions that are 3-15cm long. Mature lesions become tan with distinct dark zones of sporulation. Northern corn leaf blight can develop very rapidly, resulting in complete blighting of the leaves. There are many resistant cultivars to choose from. Cultural controls consists of deep tillage to bury debris, crop rotation, and fungicides where warranted.



## Southern corn leaf blight

Southern corn leaf blight (*Bipolaris maydis*) is not generally regarded as a serious problem as good resistance to the disease is available. However, early heavy infection in a susceptible cultivar can cause severe damage to leaves, predisposing the plant to stalk rot. Spindle shaped tan lesions with rounded ends, and buff to brown borders occasionally with a red tint, appear first on lower leaves. Race O normally attacks leaves only; whereas Race T attacks leaves, leaf sheaths, ear husks, ears, cobs, and stalks. Stalk and leaf sheath infections begin as purple spots that develop tan-gray centers. Control consists of planting resistant varieties, deep tillage to bury debris, crop rotation, and fungicides where warranted. Fungicides such as Tilt are effective against the disease.

## Northern corn leaf blight



D.G. White, APS Image Library

## Northern corn leaf blight



Sherrie Smith University of Arkansas Cooperative Extension

## Southern corn leaf blight



Sherrie Smith University of Arkansas Cooperative Extension

## Southern corn leaf blight



D.G. White, APS Image Library